

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA318617**

## **RAC1 Mouse Monoclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

**Recommend Dilution:** IHC-P (10  $\mu$ g/ml), WB (0.5 - 2  $\mu$ g/ml)

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** RAC1 antibody was raised against recombinant protein containing the full length human

Rac1.

**Formulation:** Tris-glycine, pH 7.4, 0.15 M sodium chloride, 0.05% sodium azide.

Concentration: 1 mg/ml

**Purification:** Protein G Column

Gene Name: ras-related C3 botulinum toxin substrate 1 (rho family, small GTP binding protein Rac1)

Database Link: NP 008839 Entrez Gene 19353 MouseEntrez Gene 363875 RatEntrez Gene 5879 Human

**Synonyms:** MIG5; p21-Rac1; Rac-1; TC-25

**Note:** Recognizes human Rac1 at 21kD. Slight Rac2 cross-reactivity is observed. An additional

unknown protein of ~55kD may be detected in some preparations. Species cross-reactivity:

mouse and rat.

**Protein Families:** Druggable Genome

**Protein Pathways:** Adherens junction, Amyotrophic lateral sclerosis (ALS), Axon guidance, B cell receptor

signaling pathway, Chemokine signaling pathway, Colorectal cancer, Epithelial cell signaling in

Helicobacter pylori infection, Fc epsilon RI signaling pathway, Fc gamma R-mediated


phagocytosis, Focal adhesion, Leukocyte transendothelial migration, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Pancreatic cancer, Pathways in cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, Toll-like receptor

signaling pathway, VEGF signaling pathway, Viral myocarditis, Wnt signaling pathway





## **Product images:**



Anti-RAC1 antibody IHC of human small intestine. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 10 ug/ml.