

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA319578

TIMP4 Mouse Monoclonal Antibody [Clone ID: 9:4-7]

Product data:

Product Type: Primary Antibodies

Clone Name: 9:4-7
Applications: IHC

Recommend Dilution: ELISA: 1:20,000 to 1:100,000, WB: 1:2,000 to 1:10,000, IHC: 5-10ug/mL

Reactivity: Human, Mouse

Host: Mouse

Clonality: Monoclonal

Immunogen: Anti-TIMP4 (MOUSE) Monoclonal Antibody was produced in mouse by repeated

immunizations with full legnth human TIMP4 protein followed by hybridoma development.

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Concentration: 1 mg/ml

Gene Name: TIMP metallopeptidase inhibitor 4

Database Link: NP 003247 Entrez Gene 110595 MouseEntrez Gene 7079 Human

Synonyms: TIMP metallopeptidase inhibitor 4; tissue inhibitor of metalloproteinase 4

Note: Tissue inhibitor of metalloproteinase-4 (TIMP-4) is member of the TIMP family of inhibitors of

matrix metalloproteinase (MMP) activities. In addition, TIMPs have non-MMP associated functions, including roles in growth promotion, apoptosis, and angiogenesis. All four

members of the TIMP gene family form a similar secondary structure due to the formation of six highly conserved disulfide bonds. It has been reported that some TIMPs, TIMP4 included, can bind a tetraspanin in the plasma membrane, initiating an intra-cellular signaling cascade. Elevated levels of TIMP4 have been associated with breast cancer progression and heart

disease.

Protein Families: Druggable Genome, Secreted Protein





Product images:



Immunohistochemistry of Mouse Anti-TIMP4
Antibody. Tissue: human breast carcinoma.
Fixation: formalin fixed paraffin embedded.
Antigen retrieval: not required. Primary antibody:
a) isotype specific IgG2bk monoclonal; b) TIMP-4
antibody (clone 9:4-7); and c) TIMP-4 each at 10
ug/mL. Secondary antibody: Peroxidase mouse
secondary antibody at 1:10,000 for 45 min at RT.
Localization: TIMP-4 is secreted. Staining: TIMP-4
as precipitated brown signal.