

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA320351

ICAM2 Mouse Monoclonal Antibody [Clone ID: CBRIC2/2]

Product data:

Product Type: Primary Antibodies

Clone Name: CBRIC2/2

Applications: FC

Recommend Dilution: Flow, IHC, Functional Assay, IP, WB

Reactivity: Human Host: Mouse

Clonality: Monoclonal

Formulation: Aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Concentration: 0.5mg/ml

Purification: Affinity purified

Gene Name: intercellular adhesion molecule 2

Database Link: NP 000864 Entrez Gene 3384 Human

Background: The CBRIC2/2 monoclonal antibody reacts with human ICAM-2 (CD102). ICAM-2 is a 55-65 kD

transmembrane glycoprotein possessing 2 extracellular Ig domains, a single transmembrane domain, and a short 26-amino acid cytoplasmic domain. ICAM-2 is expressed on most leukocytes, and is strongly expressed on vascular endothelial cells. ICAM-2 is a ligand of integrin CD11a/CD18 (LFA-1), and also has been reported to bind integrin CD11b/CD18 (Mac-1). ICAM-2 is an intercellular adhesion molecule involved in lymphocyte homing to sites of

inflammation. The CBRIC2/2 antibody has been reported to have blocking activity.


Synonyms: CD102

Protein Families: ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Natural killer cell mediated cytotoxicity



Product images:



Staining of normal human peripheral blood cells with 0.06 ug of Mouse IgG2a kappa Isotype Control Purified (open histogram) or 0.06 ug of Anti-Human CD102 (ICAM-2) Purified (filled histogram) followed by F (ab')2 Anti-Mouse IgG PE. Cells in the monocyte gate were used for analysis.