

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA337160**

**EPS8 Mouse Monoclonal Antibody [Clone ID: 18F5B11]** 

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 18F5B11
Applications: IHC, WB

**Recommend Dilution:** WB: 1-5ug/ml~, IHC-P: 5ug/ml

Reactivity: Human Host: Mouse

**Isotype:** IgG3, kappa **Clonality:** Monoclonal

Immunogen: Amino acids 701-882 of human EPS8 protein were used as the immunogen for the EPS8

antibody.

Formulation: PBS containing 0.05% BSA, 0.05% Sodium Azide. Store at 4C short term. Aliquot and store at -

20C long term. Avoid freeze-thaw cycles.

**Concentration:** 0.5 mg/ml

**Purification:** Protein G purified

**Gene Name:** epidermal growth factor receptor pathway substrate 8

Database Link: NP 004438 Entrez Gene 2059 Human

**Background:** EPS8, an acronym for epidermal growth factor receptor pathways substrate 8, is a signaling

protein and part of the molecular complex controlling actin reorganization (1, 2). EPS8 also plays an important role in innate immunity. Through LPS activation of TLR4 and the co

adaptor MyD88 protein, macrophages induce enhanced EPS8 activity typified by

phagocytosis- a key innate anti-microbial defense response (3). EPS8 colocalizes with TLR4 in LPS activated macrophage immunocomplexes (3). Interestingly, EPS8 also is over expressed in malignancies including breast cancers, pancreatic cancers, and colon cancers, as examples (2-4). Antibody to EPS8 provides a tool for characterizing its functional roles in motility,

adhesion, signaling pathways such as non-canonical Wnt signaling and expression in normal

and neoplastic cells (5).

Synonyms: DFNB102




## EPS8 Mouse Monoclonal Antibody [Clone ID: 18F5B11] - TA337160

**Note:** Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM sodium


citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.

**Protein Families:** Druggable Genome

## **Product images:**



Western Blot: EPS8 Antibody (18F5B11) TA337160 - analysis of EPS8 partial recombinant protein using EPS8 antibody at 1 ug/ml.



Immunohistochemistry-Paraffin: EPS8 Antibody (18F5B11) TA337160 - Formalin-fixed, paraffinembedded human salivary gland stained with EPS8 antibody (5 ug/ml), peroxidase-conjugate and DAB chromogen.