

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA500685**

## BCKDH kinase (BCKDK) Mouse Monoclonal Antibody [Clone ID: OTI11C9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI11C9
Applications: IF, IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:50, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BCKDK (NP\_005872) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.8 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 46.4 kDa

**Gene Name:** branched chain keto acid dehydrogenase kinase

Database Link: NP 005872 Entrez Gene 10295 Human

**Background:** Catalyzes the phosphorylation and inactivation of the branched-chain alpha-ketoacid

dehydrogenase complex,the key regulatory enzyme of the valine, leucine and isoleucine catabolic pathways. Key enzyme that regulate the activity state of the BCKD complex .


Synonyms: BCKDKD; BDK

**Protein Families:** Druggable Genome, Protein Kinase







### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BCKDK ([RC203601], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BCKDK. Positive lysates [LY401780] (100ug) and [LC401780] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human Ovary tissue within the normal limits using anti-BCKDK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500685)



Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-BCKDK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500685)





Anti-BCKDK mouse monoclonal antibody (TA500685) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BCKDK ([RC203601]).