

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA501563

## PRPSAP2 Mouse Monoclonal Antibody [Clone ID: OTI1E3]

### **Product data:**


| Product Type:              | Primary Antibodies   |
|----------------------------|--|
| Clone Name:                | OTI1E3   |
| Applications:              | FC, IF, IHC, WB  |
| <b>Recommend Dilution:</b> | WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100   |
| Reactivity:                | Human  |
| Host:                      | Mouse  |
| lsotype:                   | lgG1   |
| Clonality:                 | Monoclonal   |
| Immunogen:                 | Full length human recombinant protein of human PRPSAP2 (NP_002758) produced in<br>HEK293T cell.  |
| Formulation:               | PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.   |
| Concentration:             | 0.7 mg/ml  |
| Purification:              | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography<br>(protein A/G)   |
| Predicted Protein Size:    | 40.7 kDa   |
| Gene Name:                 | phosphoribosyl pyrophosphate synthetase associated protein 2   |
| Database Link:             | <u>NP_002758 Entrez Gene 5636 Human</u>  |
| Background:                | The enzyme phosphoribosylpyrophosphate synthetase (PRS) catalyzes the formation of phosphoribosylpyrophosphate which is a substrate for synthesis of purine and pyrimidine nucleotides, histidine, tryptophan and NAD. PRS exists as a complex with two catalytic subunits and two associated subunits. This gene encodes a non-catalytic associated subunit of PRS. [provided by RefSeq]. COMPLETENESS: complete on the 3' end. |
| Synonyms:                  | PAP41  |
| Protein Families:          | Druggable Genome   |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



#### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PRPSAP2 ([RC210719], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PRPSAP2. Positive lysates [LY400988] (100ug) and [LC400988] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-PRPSAP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501563)


Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-PRPSAP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501563)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-PRPSAP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501563)





HEK293T cells transfected with either [RC210719] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PRPSAP2 antibody (TA501563), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US