

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA502455

POLR2J2 Mouse Monoclonal Antibody [Clone ID: OTI6E9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6E9
Applications: FC, WB

Recommend Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human POLR2J2 (NP_116581) produced in

HEK293T cell.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.94 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 12.9 kDa

Gene Name: RNA polymerase II subunit |2

Database Link: NP 116581 Entrez Gene 246721 Human

Background: This gene is a member of the RNA polymerase II subunit 11 gene family, which includes three

genes in a cluster on chromosome 7q22.1 and a pseudogene on chromosome 7p13. The founding member of this family, DNA directed RNA polymerase II polypeptide J, has been

shown to encode a subunit of RNA polymerase II, the polymerase responsible for

synthesizing messenger RNA in eukaryotes. This locus produces multiple, alternatively spliced

transcripts that potentially express isoforms with distinct C-termini compared to DNA

directed RNA polymerase II polypeptide J. Most or all variants are spliced to include additional non-coding exons at the 3' end which makes them candidates for nonsense-mediated decay (NMD). Consequently, it is not known if this locus expresses a protein or proteins in vivo.

[provided by RefSeq]

Synonyms: HRPB11B; RPB11b1






Protein Families: Transcription Factors


Protein Pathways: Huntington's disease, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA

polymerase

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY POLR2J2 ([RC224755], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-POLR2J2. Positive lysates [LY409843] (100ug) and [LC409843] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC224755] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-POLR2J2 antibody (TA502455), and then analyzed by flow cytometry.