

## Product datasheet for **TA502736**

### **STAT5A Mouse Monoclonal Antibody [Clone ID: OTI9F7]**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	OTI9F7
<b>Applications:</b>	FC, IF, IHC, WB
<b>Recommend Dilution:</b>	WB 1:500~2000, IHC 1:150, IF 1:100, FLOW 1:100
<b>Reactivity:</b>	Human, Monkey, Rat, Dog
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG1
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Full length human recombinant protein of human STAT5A (NP_003143) produced in HEK293T cell.
<b>Formulation:</b>	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
<b>Concentration:</b>	0.73 mg/ml
<b>Purification:</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
<b>Predicted Protein Size:</b>	90.5 kDa
<b>Gene Name:</b>	signal transducer and activator of transcription 5A
<b>Database Link:</b>	<a href="#">NP_003143</a> <a href="#">Entrez Gene 24918</a> <a href="#">RatEntrez Gene 490968</a> <a href="#">DogEntrez Gene 574129</a> <a href="#">MonkeyEntrez Gene 6776</a> <a href="#">Human</a>
<b>Background:</b>	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. [provided by RefSeq]




[View online »](#)

**Synonyms:** MGF; STAT5


**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Stem cell relevant signaling - DSL/Notch pathway, Stem cell relevant signaling - JAK/STAT signaling pathway, Transcription Factors

**Protein Pathways:** Acute myeloid leukemia, Chronic myeloid leukemia, ErbB signaling pathway, Jak-STAT signaling pathway, Pathways in cancer


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY STAT5A ([RC205753], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-STAT5A. Positive lysates [LY418866] (100ug) and [LC418866] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-STAT5A monoclonal antibody.




Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-STAT5A monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-STAT5A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502736)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-STAT5A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502736)




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-STAT5A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502736)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-STAT5A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502736)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-STAT5A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502736)




Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-STAT5A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502736)




Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-STAT5A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502736)




Anti-STAT5A mouse monoclonal antibody (TA502736) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY STAT5A ([RC205753]).



HEK293T cells transfected with either [RC205753] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-STAT5A antibody (TA502736), and then analyzed by flow cytometry.



Flow cytometric Analysis of HeLa cells, using anti-STAT5A antibody (TA502736), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-STAT5A antibody (TA502736), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).