

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA502948**

## LGR5 Mouse Monoclonal Antibody [Clone ID: OTI2H7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2H7

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:500, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 250-550 of human

LGR5 (NP\_003658) produced in 293T Cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.72 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 99.8 kDa

**Gene Name:** leucine rich repeat containing G protein-coupled receptor 5


Database Link: NP 003658 Entrez Gene 8549 Human
Synonyms: FEX; GPR49; GPR67; GRP49; HG38

**Protein Families:** Druggable Genome, GPCR, Transmembrane






# **Product images:**




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-LGR5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502948)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-LGR5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502948)






Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-LGR5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502948)




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-LGR5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502948)




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY LGR5 ([RC212825], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-LGR5. Positive lysates [LY401213] (100ug) and [LC401213] (20ug) can be purchased separately from OriGene.






Immunofluorescent staining of HT29 cells using anti-LGR5 mouse monoclonal antibody (TA502948, green). Actin filaments were labeled with TRITC-phalloidin (red), and nuclear with DAPI (blue).



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-LGR5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502948)



Flow cytometric analysis of living 293T cells transfected with LGR5 overexpression plasmid ([RC212825]), Red) using anti-LGR5 antibody (TA502948). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).