

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA502991**

## **SOCS3 Mouse Monoclonal Antibody [Clone ID: OTI3D3]**

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI3D3

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SOCS3 (NP\_003946) produced in HEK293T

cell

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 24.6 kDa

Gene Name: suppressor of cytokine signaling 3

Database Link: NP 003946 Entrez Gene 9021 Human

Background: This gene encodes a member of the STAT-induced STAT inhibitor (SSI), also known as

suppressor of cytokine signaling (SOCS), family. SSI family members are cytokine-inducible negative regulators of cytokine signaling. The expression of this gene is induced by various cytokines, including IL6, IL10, and interferon (IFN)-gamma. The protein encoded by this gene

can bind to JAK2 kinase, and inhibit the activity of JAK2 kinase. Studies of the mouse counterpart of this gene suggested the roles of this gene in the negative regulation of fetal

liver hematopolesis, and placental development. [provided by RefSeq]

Synonyms: ATOD4; CIS3; Cish3; SOCS-3; SSI-3; SSI3

**Protein Families:** Druggable Genome


Protein Pathways: Adipocytokine signaling pathway, Insulin signaling pathway, Jak-STAT signaling pathway, Type

II diabetes mellitus, Ubiquitin mediated proteolysis







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SOCS3 ([RC209305], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SOCS3.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-SOCS3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502991)




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-SOCS3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502991)






Anti-SOCS3 mouse monoclonal antibody (TA502991) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SOCS3 ([RC209305] ).



Flow cytometric Analysis of Hela cells, using anti-SOCS3 antibody (TA502991), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-SOCS3 antibody (TA502991), (Red), compared to a nonspecific negative control antibody, (Blue).