

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA504271**

## **RFXANK Mouse Monoclonal Antibody [Clone ID: OTI3E7]**

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI3E7

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human RFXANK(NP\_604389) produced in HEK293T

cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 25.4 kDa

**Gene Name:** regulatory factor X associated ankyrin containing protein

Database Link: NP 604389 Entrez Gene 8625 Human

**Background:** Major histocompatibility (MHC) class II molecules are transmembrane proteins that have a

central role in development and control of the immune system. The protein encoded by this gene, along with regulatory factor X-associated protein and regulatory factor-5, forms a complex that binds to the X box motif of certain MHC class II gene promoters and activates their transcription. Once bound to the promoter, this complex associates with the non-DNA-binding factor MHC class II transactivator, which controls the cell type specificity and inducibility of MHC class II gene expression. This protein contains ankyrin repeats involved in

protein-protein interactions. Mutations in this gene have been linked to bare lymphocyte syndrome type II, complementation group B. Two transcript variants encoding different isoforms have been described for this gene, with only one isoform showing activation activity.

[provided by RefSeq]




#### RFXANK Mouse Monoclonal Antibody [Clone ID: OTI3E7] - TA504271

Synonyms: ANKRA1; BLS; F14150\_1; RFX-B


**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Antigen processing and presentation, Primary immunodeficiency


### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RFXANK ([RC223081], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RFXANK. Positive lysates [LY408725] (100ug) and [LC408725] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-RFXANK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504271)




Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-RFXANK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504271)






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-RFXANK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504271)




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-RFXANK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504271)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-RFXANK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504271)







Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-RFXANK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504271)




Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-RFXANK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504271)



Anti-RFXANK mouse monoclonal antibody (TA504271) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY RFXANK ([RC223081]).



Flow cytometric Analysis of Hela cells, using anti-RFXANK antibody (TA504271), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-RFXANK antibody (TA504271), (Red), compared to a nonspecific negative control antibody, (Blue).