

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA505177**

## **RIOK2 Mouse Monoclonal Antibody [Clone ID: OTI5D11]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5D11

**Applications:** FC, IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 277-552 of human

RIOK2(NP\_060813) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 63.1 kDa

**Gene Name:** RIO kinase 2

Database Link: NP 060813 Entrez Gene 55781 Human


Synonyms: RIO2

**Protein Families:** Druggable Genome, Protein Kinase






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RIOK2 ([RC201484], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RIOK2. Positive lysates [LY402668] (100ug) and [LC402668] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-RIOK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505177)







Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-RIOK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505177)



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-RIOK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505177)



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-RIOK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505177)



HEK293T cells transfected with either [RC201484] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-RIOK2 antibody (TA505177), and then analyzed by flow cytometry.