

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA506734

# IFNGR2 Mouse Monoclonal Antibody [Clone ID: OTI1C2]

## **Product data:**


Product Type:	Primary Antibodies
Clone Name:	OTI1C2
Applications:	IHC, WB
<b>Recommend Dilution:</b>	WB 1:500, IHC 1:150
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human IFNGR2(NP_005525) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	35 kDa
Gene Name:	interferon gamma receptor 2
Database Link:	<u>NP_005525 Entrez Gene 3460 Human</u>
Background:	This gene (IFNGR2) encodes the non-ligand-binding beta chain of the gamma interferon receptor. Human interferon-gamma receptor is a heterodimer of IFNGR1 and IFNGR2. Defects in IFNGR2 are a cause of mendelian susceptibility to mycobacterial disease (MSMD), also known as familial disseminated atypical mycobacterial infection. MSMD is a genetically heterogeneous disease with autosomal recessive, autosomal dominant or X-linked inheritance. [provided by RefSeq, Jul 2008]
Synonyms:	AF-1; IFGR2; IFNGT1; IMD28
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway, Natural killer cell mediated cytotoxicity




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IFNGR2 ([RC200674], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IFNGR2. Positive lysates [LY401697] (100ug) and [LC401697] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-IFNGR2 mouse monoclonal antibody (TA506734) at 1:150 dilution.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US