

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA507052

CHAC1 Mouse Monoclonal Antibody [Clone ID: OTI1E2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1E2

Applications: IF, IHC, WB

Recommend Dilution: WB 1:4000, IHC 1:150, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CHAC1(NP_077016) produced in HEK293T

cell.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 24.2 kDa


Gene Name: ChaC glutathione specific gamma-glutamylcyclotransferase 1

Database Link: NP 077016 Entrez Gene 79094 Human


Synonyms: MGC4504



Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CHAC1 ([RC200912], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CHAC1. Positive lysates [LY432159] (100ug) and [LC432159] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)






Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)





Immunohistochemical staining of paraffinembedded Human bladder tissue within the normal limits using anti-CHAC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA507052)



Anti-CHAC1 mouse monoclonal antibody (TA507052) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CHAC1 ([RC200912]).