

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA800258**

## MIPEP Mouse Monoclonal Antibody [Clone ID: OTI3F8]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI3F8

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 174-516 of human

MIPEP (NP\_005923) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 80.4 kDa

Gene Name: mitochondrial intermediate peptidase

Database Link: NP 005923 Entrez Gene 4285 Human

**Background:** The product of this gene performs the final step in processing a specific class of nuclear-

encoded proteins targeted to the mitochondrial matrix or inner membrane. This protein is primarily involved in the maturation of oxidative phosphorylation (OXPHOS)-related proteins. This gene may contribute to the functional effects of frataxin deficiency and the clinical

manifestations of Friedreich ataxia. [provided by RefSeq]. COMPLETENESS: complete on the 3'

end.


Synonyms: HMIP; MIP

**Protein Families:** Druggable Genome, Protease







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MIPEP ([RC202465], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MIPEP. Positive lysates [LY416973] (100ug) and [LC416973] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)






Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)






Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)






Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-MIPEP mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800258)



Anti-MIPEP mouse monoclonal antibody (TA800258) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MIPEP ([RC202465]).



HEK293T cells transfected with either [RC202465] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-MIPEP antibody (TA800258), and then analyzed by flow cytometry.