

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA800495**

### S100 beta (S100B) Mouse Monoclonal Antibody [Clone ID: OTI6C5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6C5

Applications: WB

Recommend Dilution: WB 1:2000

Reactivity: Human Host: Mouse

**Isotype:** IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human S100B (NP\_006263) produced in HEK293T

cell

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 10.5 kDa

**Gene Name:** S100 calcium binding protein B

Database Link: NP 006263 Entrez Gene 6285 Human

Background: The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-

hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21; however, this gene is located at 21q22.3. This protein may function in Neurite extension, proliferation of melanoma cells, stimulation of

Ca2+ fluxes, inhibition of PKC-mediated phosphorylation, astrocytosis and axonal

proliferation, and inhibition of microtubule assembly. Chromosomal rearrangements and altered expression of this gene have been implicated in several neurological, neoplastic, and

other types of diseases, including Alzheimer's disease, Down's syndrome, epilepsy,


amyotrophic lateral sclerosis, melanoma, and type I diabetes. [provided by RefSeq, Jul 2008]





Synonyms: NEF; S100; S100-B; S100beta

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY S100B ([RC200277], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-S100B. Positive lysates [LY416735] (100ug) and [LC416735] (20ug) can be purchased separately from OriGene.