

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA802443

## VEGF Receptor 2 (KDR) Mouse Monoclonal Antibody [Clone ID: OTI2A1]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI2A1
Applications:	WB
<b>Recommend Dilution:</b>	WB 1:2000
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 786-1356 of human KDR (NP_002244) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Gene Name:	kinase insert domain receptor
Database Link:	<u>NP_002244 Entrez Gene 3791 Human</u>
Background:	Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc Mutations of this gene are implicated in infantile capillary hemangiomas. [provided by RefSeq, May 2009]
Synonyms:	CD309; FLK1; VEGFR; VEGFR2
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Endocytosis, Focal adhesion, VEGF signaling pathway



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **Product images:**

170 — 130 — 100 — 70 — 55 — 40 — 35 — 25 — 15 — 10 —

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KDR ([RC219851], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KDR. (1:50. Positive lysates [LY400826] (100ug) and [LC400826] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US