

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA803474**

### **PPIH Mouse Monoclonal Antibody [Clone ID: OTI1C6]**

#### **Product data:**

**Applications:** 

**Product Type: Primary Antibodies** 

WB

Clone Name: OTI1C6

WB 1:2000

**Recommend Dilution:** Reactivity: Human Host: Mouse Isotype: lgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PPIH (NP\_006338) produced in E.coli.

PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. Formulation:

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 19 kDa

Gene Name: peptidylprolyl isomerase H

Database Link: NP 006338 Entrez Gene 10465 Human

Background: The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase

> (PPlase) family. PPlases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is a specific component of the complex that includes pre-mRNA processing factors PRPF3, PRPF4, and PRPF18, as well as U4/U5/U6 tri-snRNP. This protein has been shown to possess PPlase activity and may act as a protein chaperone that mediates the interactions between different proteins inside the

spliceosome. [provided by RefSeq, Jul 2008]


Synonyms: CYP-20; CYPH; SnuCyp-20; USA-CYP

**Protein Pathways:** Spliceosome






### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PPIH ([RC201529], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PPIH. Positive lysates [LY416702] (100ug) and [LC416702] (20ug) can be purchased separately from OriGene.



Western blot analysis of A549 cell lysate (35ug) by using anti-PPIH monoclonal antibody. Dilution: 1:500