

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA803670**

### MDA5 (IFIH1) Mouse Monoclonal Antibody [Clone ID: OTI11C11]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI11C11

Applications: WB

Reactivity: WB 1:2000

Human

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IFIH1 (NP\_071451) produced in HEK293T

cell

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Gene Name:** interferon induced with helicase C domain 1

Database Link: NP 071451 Entrez Gene 64135 Human

Background: DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are

putative RNA helicases. They are implicated in a number of cellular processes involving

alteration of RNA secondary structure such as translation initiation, nuclear and

mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution

patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein that is upregulated in response to treatment with beta-interferon and a protein kinase C-activating compound, mezerein. Irreversible reprogramming of melanomas can be achieved by treatment with both these agents; treatment with either agent alone only achieves

reversible differentiation. Genetic variation in this gene is associated with diabetes mellitus

insulin-dependent type 19. [provided by RefSeq, Jul 2012]


Synonyms: AGS7; Hlcd; IDDM19; MDA-5; MDA5; RLR-2





**Protein Pathways:** RIG-I-like receptor signaling pathway

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IFIH1 ([RC215661], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IFIH1. Positive lysates [LY411723] (100ug) and [LC411723] (20ug) can be purchased separately from OriGene.