

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA808539

EPLIN (LIMA1) Mouse Monoclonal Antibody [Clone ID: OTI6F7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6F7
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-275 of human

LIMA1(NP_057441) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 85 kDa

Gene Name: LIM domain and actin binding 1

Database Link: NP 057441 Entrez Gene 51474 Human

Background: This gene encodes a cytoskeleton-associated protein that inhibits actin filament

depolymerization and cross-links filaments in bundles. It is downregulated in some cancer cell lines. Alternatively spliced transcript variants encoding different isoforms have been described for this gene, and expression of some of the variants maybe independently


regulated. [provided by RefSeq, Aug 2011]

Synonyms: EPLIN; SREBP3






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY LIMA1 ([RC211066], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-LIMA1 (1:2000). Positive lysates [LY402544] (100ug) and [LC402544] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-LIMA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808539) (1:150)






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-LIMA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808539) (1:150)




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-LIMA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808539) (1:150)



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-LIMA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808539) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-LIMA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808539) (1:150)