

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA809257

PATZ1 Mouse Monoclonal Antibody [Clone ID: OTI2H4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2H4

Applications: WB

Recommend Dilution: WB 1:2000

Reactivity: Human

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PATZ1(NP_114440) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 57.4 kDa

Gene Name: POZ/BTB and AT hook containing zinc finger 1

Database Link: NP 114440 Entrez Gene 23598 Human

Background: The protein encoded by this gene contains an A-T hook DNA binding motif which usually

binds to other DNA binding structures to play an important role in chromatin modeling and transcription regulation. Its Poz domain is thought to function as a site for protein-protein interaction and is required for transcriptional repression, and the zinc-fingers comprise the DNA binding domain. Since the encoded protein has typical features of a transcription factor, it is postulated to be a repressor of gene expression. In small round cell sarcoma, this gene is fused to EWS by a small inversion of 22q, then the hybrid is thought to be translocated (t(1;22)(p36.1;q12). The rearrangement of chromosome 22 involves intron 8 of EWS and exon 1 of this gene creating a chimeric sequence containing the transactivation domain of EWS

fused to zinc finger domain of this protein. This is a distinct example of an intrachromosomal rearrangement of chromosome 22. Four alternatively spliced transcript

variants are described for this gene. [provided by RefSeq, Jul 2008]






Synonyms: dJ400N23; MAZR; PATZ; RIAZ; ZBTB19; ZNF278; ZSG

Protein Families: Transcription Factors

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PATZ1 ([RC205802], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PATZ1 (1:2000). Positive lysates [LY410365] (100ug) and [LC410365] (20ug) can be purchased separately from OriGene.